


**Serum C8 By Immunoglobulin A (Serum) Levels
In Participants ≥ 7 And < 10 Years Of Age**
C8 (ng-mL)

IgA (Serum)	Gender	N	Mean
LOW	Female	12	47.6667
	Male	13	77.7231
	Total	25	63.2960
NORMAL	Female	900	73.5884
	Male	926	79.5631
	Total	1826	76.6183
HIGH	Female	7	45.5143
	Male	11	46.3818
	Total	18	46.0444
Total	Female	919	73.0361
	Male	950	79.1537
	Total	1869	76.1456

**Serum C8 By Immunoglobulin A (Serum) Levels
In Participants ≥ 7 And < 10 Years Of Age**



Low < 34 , Normal 34-305, High > 305 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/sc012600.htm>

¹ Note, very small sample size.

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.