


**Serum C8 By Immunoglobulin A (Serum) Levels
In Participants ≥ 10 And < 12 Years Of Age**
C8 (ng-mL)

IgA (Serum)	Gender	N	Mean
LOW	Female	33	43.6061
	Male	37	78.7676
	Total	70	62.1914
NORMAL	Female	597	69.2704
	Male	654	76.2061
	Total	1251	72.8962
HIGH	Female	81	60.6975
	Male	93	77.3688
	Total	174	69.6080
Total	Female	711	67.1025
	Male	784	76.4649
	Total	1495	72.0123

**Serum C8 By Immunoglobulin A (Serum) Levels
In Participants ≥ 10 And < 12 Years Of Age**



Low < 53 , Normal 53-204, High > 204 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/sc012600.htm>

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.