


**Serum C8 By Immunoglobulin M (Serum) Levels
In Participants ≥ 7 And < 10 Years Of Age**
C8 (ng-mL)

IgM (Serum)	Gender	N	Mean
LOW	Female	7	66.0429
	Male	19	69.6421
	Total	26	68.6731
NORMAL	Female	853	74.4011
	Male	913	78.5551
	Total	1766	76.5486
HIGH	Female	59	54.1322
	Male	18	119.5556
	Total	77	69.4260
Total	Female	919	73.0361
	Male	950	79.1537
	Total	1869	76.1456

**Serum C8 By Immunoglobulin M (Serum) Levels
In Participants ≥ 7 And < 10 Years Of Age**



Low < 31 , Normal 31-208, High > 208 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/sc013100.htm>

¹ Note, very small sample size.

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.