


Serum C8 By Uric Acid (Serum) Levels In Participants ≥ 18 And < 60 Years Of Age

C8 (ng-mL)

Uric Acid (Serum)	Gender	N	Mean
LOW	Female	931	69.7944
	Male	909	85.2643
	Total	1840	77.4368
NORMAL	Female	20935	59.7651
	Male	19145	98.8166
	Total	40080	78.4188
HIGH	Female	1898	71.7616
	Male	1087	101.4050
	Total	2985	82.5563
Total	Female	23764	61.1161
	Male	21141	98.3670
	Total	44905	78.6536

Serum C8 By Uric Acid (Serum) Levels In Participants ≥ 18 And < 60 Years Of Age



Females: Low < 2.9 , Normal $2.9-6.5$, High > 6.5 (Units: mg/dL)
Males: Low < 4.3 , Normal $4.3-8.6$, High > 8.6 (Units: mg/dL)

Source: <http://www.hosp.uky.edu/ClinLab/report.pdf>

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.