


***Self-Reported Diagnosis of Larynx Cancer in Adults
Stratified by Gender and Age***

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of Larynx Cancer	Population in Quartile	Num of Reported	Percent in Quartile
Female	1	30-50	No	3012	3010	99.93%
Female	1	30-50	Yes	3012	2	0.07%
Female	1	51-older	No	2738	2738	100.0%
Female	2	30-50	No	3036	3035	99.97%
Female	2	30-50	Yes	3036	1	0.03%
Female	2	51-older	No	2749	2743	99.78%
Female	2	51-older	Yes	2749	6	0.22%
Female	3	30-50	No	3015	3015	100.0%
Female	3	51-older	No	2737	2734	99.89%
Female	3	51-older	Yes	2737	3	0.11%
Female	4	30-50	No	3022	3022	100.0%
Female	4	51-older	No	2745	2745	100.0%
Male	1	30-50	No	2601	2601	100.0%
Male	1	51-older	No	2667	2661	99.78%
Male	1	51-older	Yes	2667	6	0.22%
Male	2	30-50	No	2621	2621	100.0%
Male	2	51-older	No	2649	2643	99.77%
Male	2	51-older	Yes	2649	6	0.23%
Male	3	30-50	No	2603	2603	100.0%
Male	3	51-older	No	2662	2658	99.85%
Male	3	51-older	Yes	2662	4	0.15%
Male	4	30-50	No	2608	2607	99.96%
Male	4	30-50	Yes	2608	1	0.04%
Male	4	51-older	No	2659	2651	99.70%
Male	4	51-older	Yes	2659	8	0.30%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Larynx Cancer and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of Larynx Cancer in Female Adults Stratified by Age



Self- Reported Diagnosis of Larynx Cancer in Male Adults Stratified by Age

