


***Self-Reported Diagnosis of Liver Cancer in Adults
Stratified by Gender and Age***

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of Liver Cancer	Population in Quartile	Num of Reported	Percent in Quartile
Female	1	30-50	No	3012	3012	100.0%
Female	1	51-older	No	2734	2734	100.0%
Female	2	30-50	No	3034	3033	99.97%
Female	2	30-50	Yes	3034	1	0.03%
Female	2	51-older	No	2747	2747	100.0%
Female	3	30-50	No	3015	3015	100.0%
Female	3	51-older	No	2736	2735	99.96%
Female	3	51-older	Yes	2736	1	0.04%
Female	4	30-50	No	3022	3022	100.0%
Female	4	51-older	No	2743	2743	100.0%
Male	1	30-50	No	2601	2600	99.96%
Male	1	30-50	Yes	2601	1	0.04%
Male	1	51-older	No	2664	2663	99.96%
Male	1	51-older	Yes	2664	1	0.04%
Male	2	30-50	No	2609	2609	100.0%
Male	2	51-older	No	2642	2640	99.92%
Male	2	51-older	Yes	2642	2	0.08%
Male	3	30-50	No	2614	2614	100.0%
Male	3	51-older	No	2660	2660	100.0%
Male	4	30-50	No	2608	2608	100.0%
Male	4	51-older	No	2657	2657	100.0%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Liver Cancer and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of Liver Cancer in Female Adults Stratified by Age



Self- Reported Diagnosis of Liver Cancer in Male Adults Stratified by Age

