

***Self-Reported Diagnosis of Prostate Cancer in Male Adults  
Stratified by Age***

<b>Quartile of C8 PFOA</b>	<b>AGE GROUP</b>	<b>Diagnosis of Prostate Cancer</b>	<b>Population in Quartile</b>	<b>Num of Reported</b>	<b>Percent in Quartile</b>
1	30-50	No	2601	2601	100.0%
1	51-older	No	2663	2572	96.58%
1	51-older	Yes	2663	91	3.42%
2	30-50	No	2621	2620	99.96%
2	30-50	Yes	2621	1	0.04%
2	51-older	No	2649	2543	96.00%
2	51-older	Yes	2649	106	4.00%
3	30-50	No	2603	2602	99.96%
3	30-50	Yes	2603	1	0.04%
3	51-older	No	2658	2534	95.33%
3	51-older	Yes	2658	124	4.67%
4	30-50	No	2608	2606	99.92%
4	30-50	Yes	2608	2	0.08%
4	51-older	No	2657	2563	96.46%
4	51-older	Yes	2657	94	3.54%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Prostate Cancer and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

# Self- Reported Diagnosis of Prostate Cancer in Male Adults Stratified by Age

