

Self-Reported Clinical Symptoms: Constipation


All Age and Gender Groups

NOTE: Symptoms were self reported by participants and were not independently verified with medical records or by a health care professional. Limited information about the characteristics of the self-reported symptoms, such as severity, duration, or history, was collected. Inferences about population health or associations based on these self-reported symptoms should be made cautiously.


SELF-REPORTED SYMPTOMS BY GENDER AND AGE GROUP: CONSTIPATION

| Gender | | | | Constipation | | | | Total |
|--------|-----------|-------|-------------|--------------|-----------|--------|-------|--------|
| | | | | Frequently | Sometimes | Rarely | Never | |
| Female | Age Group | 0-10 | Count | 109 | 390 | 639 | 1552 | 2690 |
| | | | Percent (%) | 4.1% | 14.5% | 23.8% | 57.7% | 100.0% |
| | | 11-20 | Count | 159 | 568 | 1168 | 2875 | 4770 |
| | | | Percent (%) | 3.3% | 11.9% | 24.5% | 60.3% | 100.0% |
| | | 21-30 | Count | 510 | 1208 | 1689 | 2001 | 5408 |
| | | | Percent (%) | 9.4% | 22.3% | 31.2% | 37.0% | 100.0% |
| | | 31-40 | Count | 623 | 1609 | 1892 | 1667 | 5791 |
| | | | Percent (%) | 10.8% | 27.8% | 32.7% | 28.8% | 100.0% |
| | | 41-50 | Count | 778 | 1791 | 2056 | 1788 | 6413 |
| | | | Percent (%) | 12.1% | 27.9% | 32.1% | 27.9% | 100.0% |
| | | 51-60 | Count | 572 | 1553 | 1620 | 1512 | 5257 |
| | | | Percent (%) | 10.9% | 29.5% | 30.8% | 28.8% | 100.0% |
| | | 61-70 | Count | 308 | 906 | 1001 | 1114 | 3329 |
| | | | Percent (%) | 9.3% | 27.2% | 30.1% | 33.5% | 100.0% |
| | | 71-80 | Count | 192 | 370 | 418 | 591 | 1571 |
| | | | Percent (%) | 12.2% | 23.6% | 26.6% | 37.6% | 100.0% |
| | | >80 | Count | 82 | 162 | 109 | 206 | 559 |
| | | | Percent (%) | 14.7% | 29.0% | 19.5% | 36.9% | 100.0% |
| | | Total | Count | 3333 | 8557 | 10592 | 13306 | 35788 |
| | | | Percent (%) | 9.3% | 23.9% | 29.6% | 37.2% | 100.0% |
| Male | Age Group | 0-10 | Count | 92 | 338 | 630 | 1643 | 2703 |
| | | | Percent (%) | 3.4% | 12.5% | 23.3% | 60.8% | 100.0% |
| | | 11-20 | Count | 89 | 488 | 1126 | 3258 | 4961 |
| | | | Percent (%) | 1.8% | 9.8% | 22.7% | 65.7% | 100.0% |
| | | 21-30 | Count | 103 | 502 | 1197 | 2797 | 4599 |
| | | | Percent (%) | 2.2% | 10.9% | 26.0% | 60.8% | 100.0% |
| | | 31-40 | Count | 128 | 650 | 1453 | 2649 | 4880 |
| | | | Percent (%) | 2.6% | 13.3% | 29.8% | 54.3% | 100.0% |
| | | 41-50 | Count | 169 | 866 | 1759 | 2880 | 5674 |
| | | | Percent (%) | 3.0% | 15.3% | 31.0% | 50.8% | 100.0% |
| | | 51-60 | Count | 164 | 814 | 1652 | 2443 | 5073 |
| | | | Percent (%) | 3.2% | 16.0% | 32.6% | 48.2% | 100.0% |
| | | 61-70 | Count | 134 | 542 | 1123 | 1606 | 3405 |
| | | | Percent (%) | 3.9% | 15.9% | 33.0% | 47.2% | 100.0% |
| | | 71-80 | Count | 83 | 308 | 481 | 725 | 1597 |
| | | | Percent (%) | 5.2% | 19.3% | 30.1% | 45.4% | 100.0% |
| | | >80 | Count | 38 | 83 | 92 | 137 | 350 |
| | | | Percent (%) | 10.9% | 23.7% | 26.3% | 39.1% | 100.0% |
| | | Total | Count | 1000 | 4591 | 9513 | 18138 | 33242 |
| | | | Percent (%) | 3.0% | 13.8% | 28.6% | 54.6% | 100.0% |

CONSTIPATION IN FEMALES BY AGE GROUP



CONSTIPATION IN MALES BY AGE GROUP



CONSTIPATION BY GENDER AND AGE GROUP

