

SELF-REPORTED DIAGNOSIS OF LARYNX CANCER

STRATIFIED BY GENDER AND AGE-GROUP

				Larynx Cancer			Total
				NO CANCER/ NO RESPONSE	PRIMARY	SECONDARY	
Gender	Age Group						
Female	0-10	Count	2690	0	0	2690	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	11-20	Count	4770	0	0	4770	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	21-30	Count	5408	0	0	5408	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	31-40	Count	5791	0	0	5791	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	41-50	Count	6409	4	0	6413	
		Percent (%)	99.9%	.1%	.0%	100.0%	
	51-60	Count	5254	3	0	5257	
		Percent (%)	99.9%	.1%	.0%	100.0%	
	61-70	Count	3325	4	0	3329	
		Percent (%)	99.9%	.1%	.0%	100.0%	
	71-80	Count	1570	1	0	1571	
		Percent (%)	99.9%	.1%	.0%	100.0%	
	>80	Count	558	0	1	559	
Percent (%)		99.8%	.0%	.2%	100.0%		
Total		Count	35775	12	1	35788	
		Percent (%)	100.0%	.0%	.0%	100.0%	
Male	0-10	Count	2703	0	0	2703	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	11-20	Count	4961	0	0	4961	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	21-30	Count	4599	0	0	4599	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	31-40	Count	4880	0	0	4880	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	41-50	Count	5673	1	0	5674	
		Percent (%)	100.0%	.0%	.0%	100.0%	
	51-60	Count	5066	7	0	5073	
		Percent (%)	99.9%	.1%	.0%	100.0%	
	61-70	Count	3394	10	1	3405	
		Percent (%)	99.7%	.3%	.0%	100.0%	
	71-80	Count	1592	5	0	1597	
		Percent (%)	99.7%	.3%	.0%	100.0%	
	>80	Count	348	2	0	350	
Percent (%)		99.4%	.6%	.0%	100.0%		
Total		Count	33216	25	1	33242	
		Percent (%)	99.9%	.1%	.0%	100.0%	