


Serum C8 By Uric Acid (Serum) Levels In Participants ≥ 90 Years Of Age C8 (ng-mL)

Uric Acid (Serum)	Gender	N	Mean
LOW	Male	2	78.1000
	Total	2	78.1000
NORMAL	Female	46	96.7000
	Male	16	147.4438
	Total	62	109.7952
HIGH	Female	6	72.2333
	Male	2	8.0500
	Total	8	56.1875
Total	Female	52	93.8769
	Male	20	126.5700
	Total	72	102.9583

Serum C8 By Uric Acid (Serum) Levels In Participants ≥ 90 Years Of Age



Females: Low < 2.2 , Normal $2.2-7.7$, High > 7.7 (Units: mg/dL)
 Males: Low < 4.3 , Normal $4.3-8.6$, High > 8.6 (Units: mg/dL)

Source: <http://www.hosp.uky.edu/ClinLab/report.pdf>

¹ Note, very small sample size.

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.