


Serum C8 By BUN:Creatinine (Ratio) Levels In Participants <18 Years Of Age

C8 (ng-mL)

BUN:Creatinine	Gender	N	Mean
LOW	Female	110	40.0736
	Male	152	58.4007
	Total	262	50.7061
NORMAL	Female	3108	59.0153
	Male	3040	73.9022
	Total	6148	66.3764
HIGH	Female	1887	67.7128
	Male	2242	76.7434
	Total	4129	72.6163
Total	Female	5105	61.8221
	Male	5434	74.6409
	Total	10539	68.4315

Serum C8 By BUN:Creatinine (Ratio) Levels In Participants <18 Years Of Age



Low <10:1, Normal 10:1-20:1, High >20:1 (Ratio)
 Source: <http://www.med.nyu.edu/healthwise/article.html?hwid=aa36271>

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.