


Serum C8 By C-Reactive Protein (CRP) Levels In Participants ≥ 18 Years Of Age

C8 (ng-mL)

C-Reactive Protein (CRP)	Gender	N	Mean
NORMAL	Female	21117	72.0832
	Male	22532	105.7078
	Total	43649	89.4405
HIGH	Female	8264	64.2670
	Male	4124	88.4156
	Total	12388	72.3061
Total	Female	29381	69.8847
	Male	26656	103.0325
	Total	56037	85.6527

Serum C8 By C-Reactive Protein (CRP) Levels In Participants ≥ 18 Years Of Age



Normal 0-4.9, High >4.9 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/se019500.htm>

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.