


Serum C8 By Globulin (Total) Levels In Participants <18 Years Of Age

C8 (ng-mL)

Globulin (Total)	Gender	N	Mean
LOW	Female	90	66.9778
	Male	133	75.8075
	Total	223	72.2439
NORMAL	Female	5012	61.7307
	Male	5301	74.6116
	Total	10313	68.3516
HIGH	Female	3	59.7667
	Total	3	59.7667
Total	Female	5105	61.8221
	Male	5434	74.6409
	Total	10539	68.4315

Serum C8 By Globulin (Total) Levels In Participants <18 Years Of Age



Low <2.2, Normal 2.2-4.2, High >4.2 (Units: g/dL)
 Source: http://www.drstandley.com/labvalues_proteins.shtml

¹ Note, very small sample size.

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.