


**Serum C8 By Immunoglobulin A (Serum) Levels
In Participants ≥ 16 And < 20 Years Of Age**
C8 (ng-mL)

IgA (Serum)	Gender	N	Mean
LOW	Female	70	59.2829
	Male	70	59.2257
	Total	140	59.2543
NORMAL	Female	1934	48.6062
	Male	1904	67.8906
	Total	3838	58.1730
HIGH	Female	53	38.2113
	Male	66	65.3045
	Total	119	53.2378
Total	Female	2057	48.7017
	Male	2040	67.5096
	Total	4097	58.0666

**Serum C8 By Immunoglobulin A (Serum) Levels
In Participants ≥ 16 And < 20 Years Of Age**



Low < 61 , Normal 61-348, High > 348 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/sc012600.htm>

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.