


**Serum C8 By Immunoglobulin A (Serum) Levels
In Participants ≥ 20 Years Of Age**
C8 (ng-mL)

IgA (Serum)	Gender	N	Mean
LOW	Female	735	89.4137
	Male	602	125.8193
	Total	1337	105.8058
NORMAL	Female	26221	70.3014
	Male	23371	104.8063
	Total	49592	86.5624
HIGH	Female	1378	71.1735
	Male	1667	92.5522
	Total	3045	82.8774
Total	Female	28334	70.8396
	Male	25640	104.5029
	Total	53974	86.8311

**Serum C8 By Immunoglobulin A (Serum) Levels
In Participants ≥ 20 Years Of Age**



Low <70, Normal 70-400, High >400 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/sc012600.htm>

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.