


**Serum C8 By Immunoglobulin G (Serum) Levels
In Participants ≥ 1 And < 4 Years Of Age**
C8 (ng-mL)

IgG (Serum)	Gender	N	Mean
LOW	Female	9	70.5000
	Male	9	191.1556
	Total	18	130.8278
NORMAL	Female	129	80.2403
	Male	140	93.4479
	Total	269	87.1141
HIGH	Female	33	49.7970
	Male	27	41.5333
	Total	60	46.0783
Total	Female	171	73.8526
	Male	176	90.4801
	Total	347	82.2862

**Serum C8 By Immunoglobulin G (Serum) Levels
In Participants ≥ 1 And < 4 Years Of Age**



Low < 453 , Normal $453-916$, High > 916 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/sc012800.htm>

¹ Note, very small sample size.

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.