


**Serum C8 By Immunoglobulin G (Serum) Levels
In Participants ≥ 10 And < 12 Years Of Age**
C8 (ng-mL)

IgG (Serum)	Gender	N	Mean
LOW	Female	36	66.6806
	Male	56	102.0554
	Total	92	88.2130
NORMAL	Female	665	67.7418
	Male	721	74.9222
	Total	1386	71.4771
HIGH	Female	10	26.1100
	Male	7	30.6429
	Total	17	27.9765
Total	Female	711	67.1025
	Male	784	76.4649
	Total	1495	72.0123

**Serum C8 By Immunoglobulin G (Serum) Levels
In Participants ≥ 10 And < 12 Years Of Age**



Low < 698 , Normal 698-1560, High > 1560 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/sc012800.htm>

¹ Note, very small sample size.

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.