


**Serum C8 By Immunoglobulin G (Serum) Levels
In Participants ≥ 20 Years Of Age**
C8 (ng-mL)

IgG (Serum)	Gender	N	Mean
LOW	Female	1610	87.2673
	Male	1390	117.1015
	Total	3000	101.0904
NORMAL	Female	26043	69.2410
	Male	23756	103.7867
	Total	49799	85.7206
HIGH	Female	680	93.2615
	Male	494	103.4957
	Total	1174	97.5679
Total	Female	28333	70.8419
	Male	25640	104.5029
	Total	53973	86.8326

**Serum C8 By Immunoglobulin G (Serum) Levels
In Participants ≥ 20 Years Of Age**



Low <700, Normal 700-1600, High >1600 (Units: mg/dL)

Source: <http://www.labcorp.com/datasets/labcorp/html/chapter/mono/sc012800.htm>

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.