


Serum C8 By Uric Acid (Serum) Levels In Participants ≥ 10 And < 13 Years Of Age C8 (ng-mL)

Uric Acid (Serum)	Gender	N	Mean
LOW	Female	18	37.8944
	Male	8	39.3500
	Total	26	38.3423
NORMAL	Female	998	64.0907
	Male	1095	79.8751
	Total	2093	72.3486
HIGH	Female	80	55.8850
	Male	167	70.4497
	Total	247	65.7324
Total	Female	1096	63.0615
	Male	1270	78.3804
	Total	2366	71.2842

Serum C8 By Uric Acid (Serum) Levels In Participants ≥ 10 And < 13 Years Of Age



Females: Low < 2.5 , Normal 2.5-5.9, High > 5.9 (Units: mg/dL)

Males: Low < 2.2 , Normal 2.2-5.8, High > 5.8 (Units: mg/dL)

Source: <http://www.hosp.uky.edu/ClinLab/report.pdf>

¹ Note, very small sample size.

The WVU website is a communication vehicle to depict associations or their absence for public use. These tables and graphs show many comparisons between lab tests and corresponding population serum PFOA (C8) levels. When it appears that there is a clear relationship between serum C8 and a clinical laboratory value, the meaning of that relationship still requires thought and discussion. Some of the relationships, while real, are weak and not likely to be important. Several are strong, interesting and potentially important, and none of them can be taken to show an etiologic (cause and effect) relationship or its absence without more work. When it comes to causes, scientists interpret these preliminary data with deference to additional work that needs to be done.

These data concerning associations are for public use. They will receive additional collaborative work in peer review format. We hope they prompt public curiosity and suggestions of interested scientists.