

***Self-Reported Diagnosis of Uterine Fibroids in Female  
Stratified by Age***

<b>Gender</b>	<b>Quartile of C8 PFOA</b>	<b>AGE GROUP</b>	<b>Diagnosis of Uterine Fibroids</b>	<b>Population in Quartile</b>	<b>Num of Reported</b>	<b>Percent in Quartile</b>
Female	1	30-50	No	3012	2569	85.29%
Female	1	30-50	Yes	3012	443	14.71%
Female	1	51-older	No	2738	2073	75.71%
Female	1	51-older	Yes	2738	665	24.29%
Female	2	30-50	No	3036	2586	85.18%
Female	2	30-50	Yes	3036	450	14.82%
Female	2	51-older	No	2749	2122	77.19%
Female	2	51-older	Yes	2749	627	22.81%
Female	3	30-50	No	3015	2650	87.89%
Female	3	30-50	Yes	3015	365	12.11%
Female	3	51-older	No	2744	2144	78.13%
Female	3	51-older	Yes	2744	600	21.87%
Female	4	30-50	No	3022	2625	86.86%
Female	4	30-50	Yes	3022	397	13.14%
Female	4	51-older	No	2739	2132	77.84%
Female	4	51-older	Yes	2739	607	22.16%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Uterine Fibroids and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

# Self- Reported Diagnosis of Uterine Fibroids in Female Adults Stratified by Age

