


***Self-Reported Diagnosis of Stomach Cancer in Adults
Stratified by Gender and Age***

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of Stomach Cancer	Population in Quartile	Num of Reported	Percent in Quartile
Female	1	30-50	No	3011	3010	99.97%
Female	1	30-50	Yes	3011	1	0.03%
Female	1	51-older	No	2735	2732	99.89%
Female	1	51-older	Yes	2735	3	0.11%
Female	2	30-50	No	3035	3035	100.0%
Female	2	51-older	No	2747	2747	100.0%
Female	3	30-50	No	3015	3013	99.93%
Female	3	30-50	Yes	3015	2	0.07%
Female	3	51-older	No	2743	2739	99.85%
Female	3	51-older	Yes	2743	4	0.15%
Female	4	30-50	No	3022	3022	100.0%
Female	4	51-older	No	2738	2736	99.93%
Female	4	51-older	Yes	2738	2	0.07%
Male	1	30-50	No	2601	2599	99.92%
Male	1	30-50	Yes	2601	2	0.08%
Male	1	51-older	No	2668	2662	99.78%
Male	1	51-older	Yes	2668	6	0.22%
Male	2	30-50	No	2621	2620	99.96%
Male	2	30-50	Yes	2621	1	0.04%
Male	2	51-older	No	2648	2647	99.96%
Male	2	51-older	Yes	2648	1	0.04%
Male	3	30-50	No	2603	2603	100.0%
Male	3	51-older	No	2661	2660	99.96%
Male	3	51-older	Yes	2661	1	0.04%
Male	4	30-50	No	2608	2608	100.0%
Male	4	51-older	No	2659	2657	99.92%
Male	4	51-older	Yes	2659	2	0.08%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Stomach Cancer and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of Stomach Cancer in Female Adults Stratified by Age



Self- Reported Diagnosis of Stomach Cancer in Male Adults Stratified by Age

