

***Self-Reported Diagnosis of Testicular Cancer in Male Adults  
Stratified by Age***

<b>Quartile of C8 PFOA</b>	<b>Age Group</b>	<b>Diagnosis of Testicular Cancer</b>	<b>Population in Quartile</b>	<b>Num of Reported</b>	<b>Percent in Quartile</b>
1	30-50	No	2601	2596	99.81%
1	30-50	Yes	2601	5	0.19%
1	51-older	No	2668	2666	99.93%
1	51-older	Yes	2668	2	0.07%
2	30-50	No	2621	2613	99.69%
2	30-50	Yes	2621	8	0.31%
2	51-older	No	2649	2646	99.89%
2	51-older	Yes	2649	3	0.11%
3	30-50	No	2603	2597	99.77%
3	30-50	Yes	2603	6	0.23%
3	51-older	No	2662	2658	99.85%
3	51-older	Yes	2662	4	0.15%
4	30-50	No	2608	2598	99.62%
4	30-50	Yes	2608	10	0.38%
4	51-older	No	2659	2654	99.81%
4	51-older	Yes	2659	5	0.19%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Testicular Cancer and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

# *Self- Reported Diagnosis of Testicular Cancer in Male Adults Stratified by Age*

