

***Self-Reported Diagnosis of Cervix Cancer in Female Adults  
Stratified by Age***

<b>Quartile of C8 PFOA</b>	<b>AGE GROUP</b>	<b>Diagnosis of Cervix Cancer</b>	<b>Population in Quartile</b>	<b>Num of Reported</b>	<b>Percent in Quartile</b>
1	30-50	No	5638	5527	98.03%
1	30-50	Yes	5638	111	1.97%
1	51-older	No	5394	5315	98.54%
1	51-older	Yes	5394	79	1.46%
2	30-50	No	5629	5545	98.51%
2	30-50	Yes	5629	84	1.49%
2	51-older	No	5398	5333	98.80%
2	51-older	Yes	5398	65	1.20%
3	30-50	No	5618	5539	98.59%
3	30-50	Yes	5618	79	1.41%
3	51-older	No	5401	5334	98.76%
3	51-older	Yes	5401	67	1.24%
4	30-50	No	5624	5562	98.90%
4	30-50	Yes	5624	62	1.10%
4	51-older	No	5397	5346	99.06%
4	51-older	Yes	5397	51	0.94%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Cervix Cancer and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

# Self- Reported Diagnosis of Cervix Cancer in Female Adults Stratified by Age

