


***Self-Reported Diagnosis of GallBladder Cancer in Adults
Stratified by Gender and Age***

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of GallBladder Cancer	Population in Quartile	Num of Reported	Percent in Quartile
Female	1	30-50	No	3012	3012	100.0%
Female	1	51-older	No	2738	2738	100.0%
Female	2	30-50	No	3036	3036	100.0%
Female	2	51-older	No	2749	2748	99.96%
Female	2	51-older	Yes	2749	1	0.04%
Female	3	30-50	No	3015	3015	100.0%
Female	3	51-older	No	2736	2735	99.96%
Female	3	51-older	Yes	2736	1	0.04%
Female	4	30-50	No	3021	3021	100.0%
Female	4	51-older	No	2744	2743	99.96%
Female	4	51-older	Yes	2744	1	0.04%
Male	1	30-50	No	2601	2601	100.0%
Male	1	51-older	No	2668	2667	99.96%
Male	1	51-older	Yes	2668	1	0.04%
Male	2	30-50	No	2621	2621	100.0%
Male	2	51-older	No	2649	2649	100.0%
Male	3	30-50	No	2603	2603	100.0%
Male	3	51-older	No	2661	2660	99.96%
Male	3	51-older	Yes	2661	1	0.04%
Male	4	30-50	No	2608	2608	100.0%
Male	4	51-older	No	2659	2659	100.0%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported GallBladder Cancer and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of GallBladder Cancer in Female Adults Stratified by Age



Self- Reported Diagnosis of GallBladder Cancer in Male Adults Stratified by Age

