


Self-Reported Osteoarthritis and C8/PFOA

Adults (>=18YO) Only

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported osteoarthritis and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

				Self-Reported Osteoarthritis		
				No/No Response	Yes	Total
Female	Quartile Group -- C8	1	Count	8502	670	9172
			% within Quartile Group -- C8	92.7%	7.3%	100.0%
			% within Self-Reported Osteoarthritis	32.2%	21.8%	31.1%
		2	Count	6625	697	7322
			% within Quartile Group -- C8	90.5%	9.5%	100.0%
			% within Self-Reported Osteoarthritis	25.1%	22.7%	24.8%
		3	Count	5978	754	6732
			% within Quartile Group -- C8	88.8%	11.2%	100.0%
			% within Self-Reported Osteoarthritis	22.6%	24.5%	22.8%
		4	Count	5289	953	6242
			% within Quartile Group -- C8	84.7%	15.3%	100.0%
			% within Self-Reported Osteoarthritis	20.0%	31.0%	21.2%
	Total		Count	26394	3074	29468
% within Quartile Group -- C8			89.6%	10.4%	100.0%	
% within Self-Reported Osteoarthritis			100.0%	100.0%	100.0%	
Male	Quartile Group -- C8	1	Count	4618	224	4842
			% within Quartile Group -- C8	95.4%	4.6%	100.0%
			% within Self-Reported Osteoarthritis	18.2%	16.8%	18.1%
		2	Count	6580	281	6861
			% within Quartile Group -- C8	95.9%	4.1%	100.0%
			% within Self-Reported Osteoarthritis	25.9%	21.1%	25.7%
		3	Count	6766	362	7128
			% within Quartile Group -- C8	94.9%	5.1%	100.0%
			% within Self-Reported Osteoarthritis	26.7%	27.2%	26.7%
		4	Count	7413	463	7876
			% within Quartile Group -- C8	94.1%	5.9%	100.0%
			% within Self-Reported Osteoarthritis	29.2%	34.8%	29.5%
	Total		Count	25377	1330	26707
% within Quartile Group -- C8			95.0%	5.0%	100.0%	
% within Self-Reported Osteoarthritis			100.0%	100.0%	100.0%	

Participants with Self-Reported Osteoarthritis within C8/PFOA Population Quartile Group



C8/PFOA for Participants Stratified by Self-Reported Osteoarthritis

