

***Self-Reported Diagnosis of Prostatitis in Male
Stratified and Age***

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of Prostatitis	Population in Quartile	Num of Reported	Percent in Quartile
Male	1	30-50	No	2601	2588	99.50%
Male	1	30-50	Yes	2601	13	0.50%
Male	1	51-older	No	2668	2618	98.13%
Male	1	51-older	Yes	2668	50	1.87%
Male	2	30-50	No	2621	2613	99.69%
Male	2	30-50	Yes	2621	8	0.31%
Male	2	51-older	No	2649	2592	97.85%
Male	2	51-older	Yes	2649	57	2.15%
Male	3	30-50	No	2603	2592	99.58%
Male	3	30-50	Yes	2603	11	0.42%
Male	3	51-older	No	2662	2618	98.35%
Male	3	51-older	Yes	2662	44	1.65%
Male	4	30-50	No	2608	2590	99.31%
Male	4	30-50	Yes	2608	18	0.69%
Male	4	51-older	No	2659	2631	98.95%
Male	4	51-older	Yes	2659	28	1.05%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Prostatitis and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of Prostatitis in Male Adults Stratified by Age

