


Summary Results for Hematocrit for Participants 2-9 Years of Age

HEMATOCRIT IN FEMALE SUBJECTS AGED 2-9 YEARS

PARTICIPANTS=1467, MEAN=38.20, STANDARD DEVIATION=2.29

1st PERCENTILE=33.10, MEDIAN=38.20, 99th PERCENTILE=43.60




NORMAL RANGE 33-43 %, TAKEN FROM <http://www.hosp.uky.edu/ClinLab/report.pdf>

HEMATOCRIT IN MALE SUBJECTS AGED 2-9 YEARS

PARTICIPANTS=1461, MEAN=38.23, STANDARD DEVIATION=2.34


1st PERCENTILE=33.10, MEDIAN=38.20, 99th PERCENTILE=44.40



NORMAL RANGE 33-43 %, TAKEN FROM <http://www.hosp.uky.edu/ClinLab/report.pdf>

HEMATOCRIT IN SUBJECTS AGED 2-9 YEARS

BY HEMATOCRIT CATEGORY WITHIN GENDER (PARTICIPANTS=2928)



1. FOR FEMALE IN THIS AGE GROUP, NORMAL RANGE IS 33-43 %
2. FOR MALE IN THIS AGE GROUP, NORMAL RANGE IS 33-43 %
3. RANGES TAKEN FROM <http://www.hosp.uky.edu/ClinLab/report.pdf>
4. RESULTS BASED ON LAB VALUES, NOT DIAGNOSIS

Reader's Guide to Understanding This Information:

More than 95% of all children 2-9 years of age had expected levels of hematocrit.