

SELF-REPORTED DIAGNOSIS OF PANCREATIC CANCER

STRATIFIED BY GENDER AND AGE-GROUP

			Pancreatic Cancer			Total	
			NO CANCER/ NO RESPONSE	PRIMARY	SECONDARY		
Female	Age Group	0-10	Count	2690	0	0	2690
			Percent (%)	100.0%	.0%	.0%	100.0%
		11-20	Count	4770	0	0	4770
			Percent (%)	100.0%	.0%	.0%	100.0%
		21-30	Count	5408	0	0	5408
			Percent (%)	100.0%	.0%	.0%	100.0%
		31-40	Count	5791	0	0	5791
			Percent (%)	100.0%	.0%	.0%	100.0%
		41-50	Count	6413	0	0	6413
			Percent (%)	100.0%	.0%	.0%	100.0%
		51-60	Count	5257	0	0	5257
			Percent (%)	100.0%	.0%	.0%	100.0%
		61-70	Count	3328	1	0	3329
			Percent (%)	100.0%	.0%	.0%	100.0%
		71-80	Count	1570	1	0	1571
	Percent (%)		99.9%	.1%	.0%	100.0%	
	>80	Count	558	0	1	559	
		Percent (%)	99.8%	.0%	.2%	100.0%	
	Total		Count	35785	2	1	35788
			Percent (%)	100.0%	.0%	.0%	100.0%
Male	Age Group	0-10	Count	2703	0	0	2703
			Percent (%)	100.0%	.0%	.0%	100.0%
		11-20	Count	4961	0	0	4961
			Percent (%)	100.0%	.0%	.0%	100.0%
		21-30	Count	4599	0	0	4599
			Percent (%)	100.0%	.0%	.0%	100.0%
		31-40	Count	4880	0	0	4880
			Percent (%)	100.0%	.0%	.0%	100.0%
		41-50	Count	5673	0	1	5674
			Percent (%)	100.0%	.0%	.0%	100.0%
		51-60	Count	5072	1	0	5073
			Percent (%)	100.0%	.0%	.0%	100.0%
		61-70	Count	3403	2	0	3405
			Percent (%)	99.9%	.1%	.0%	100.0%
		71-80	Count	1597	0	0	1597
	Percent (%)		100.0%	.0%	.0%	100.0%	
	>80	Count	348	2	0	350	
		Percent (%)	99.4%	.6%	.0%	100.0%	
	Total		Count	33236	5	1	33242
			Percent (%)	100.0%	.0%	.0%	100.0%